

# GLOBAL CR FUNCTIONS ON PARABOLIC CR MANIFOLDS

ANDREA ALTOMANI

**ABSTRACT.** Let  $M$  be an orbit of a real semisimple Lie group  $G_0$  acting on a complex flag manifold  $G/Q$  of its complexification  $G$ . We study the space of global CR functions on  $M$  and characterize those  $M$  which are strictly locally CR separable, i.e. those for which global CR functions induce local embeddings in  $\mathbb{C}^n$ .

## INTRODUCTION

Let  $G$  be a complex connected semisimple Lie group,  $Q$  a parabolic subgroup and  $G_0$  a connected real form of  $G$ . The orbits of  $G_0$  in the flag manifold  $Z = G/Q$  are finitely many, and carry a natural CR structure induced by their embedding in  $Z$ . In [2] these  $G_0$ -orbits are called *parabolic CR manifolds*, and their CR structure is investigated by associating to them their CR algebras [6].

In this paper we describe the space of global CR functions on parabolic CR manifolds. This was done by Wolf [9] in the case of open orbits; he proved that an open orbit is holomorphically separable if and only if it is a bounded symmetric domain. For general orbits a similar statement is true, but there are some important differences from the open case.

We need to introduce two notions of local CR separability for a CR manifold  $M$ . *Weak local CR separability* means that global CR functions locally separate points, that is that there exist global CR maps that give local CR immersions into complex Euclidean spaces. The stronger condition of *strict local CR separability* instead requires the existence of global CR maps into complex Euclidean spaces which are locally *CR embeddings*. These two conditions are not equivalent, as Example 3 below shows.

We prove that every parabolic CR manifold  $M$  admits a unique fibration  $\rho: M \rightarrow M'$  over a parabolic CR manifold  $M'$  such that  $M'$  is strictly CR separable and  $\rho$  induces an isomorphism  $\rho^*: \mathcal{O}(M') \rightarrow \mathcal{O}(M)$  of the spaces of global CR functions on  $M$  and  $M'$ .

Then we characterize strictly locally CR separable parabolic CR manifolds as those for which the fiber of the fundamental CR reduction is either a bounded symmetric domain or a smooth stratum in the boundary of a bounded symmetric domain.

Finally we discuss some examples in detail.

---

*Date:* 27 February 2007.

*2000 Mathematics Subject Classification.* Primary: 32V10; Secondary: 14M15, 32C09.

*Key words and phrases.* parabolic CR manifold, real group orbit, complex flag manifold, global CR functions.

I wish to thank prof. M. Nacinovich for several useful discussions during the preparation of this paper

## 1. DEFINITIONS AND PRELIMINARY RESULTS

Let  $M = (M, HM, J)$  be a CR manifold (see e.g. [3] for the relevant definitions). We denote by  $\mathcal{O}(M)$  the space of *smooth CR functions* on  $M$ .

We say that  $M$  is (globally) *weakly CR separable* if CR functions separate points of  $M$ , that is if for every pair  $x \neq y$  of points of  $M$  there exists a CR function  $f \in \mathcal{O}(M)$  such that  $f(x) \neq f(y)$ . We also say that  $M$  is *weakly locally CR separable* if every point of  $M$  has an open neighborhood  $U$  such that global CR functions on  $M$  separate points of  $U$ .

To introduce the notion of strict CR separability of a CR manifold we need some more notation. Let  $\mathcal{S} = \mathcal{S}(M) \subset \Gamma(T^{\mathbb{C}}M)$  be the space of complex vector fields  $X$  on  $M$  such that  $X(f) = 0$  for all  $f \in \mathcal{O}(M)$ , and let  $S = S(M) \subset T^{\mathbb{C}}M$  be the vector distribution defined at  $x \in M$  by:

$$(1.1) \quad S_x = \{X_x \mid X \in \mathcal{S}\}.$$

By the definition of CR functions,  $T^{(0,1)}M \subset S$ . We say that  $M$  is *strictly locally CR separable* at a point  $x$  if  $S_x = T_x^{(0,1)}M$ , and that  $M$  is strictly locally CR separable if it is strictly locally CR separable at each point. We have:

**Lemma 1.1.** *If  $M$  is strictly locally CR separable then  $M$  is weakly locally CR separable.*

*Proof.* Assume that there exist a point  $p \in M$  and two sequences  $x_n$  and  $y_n$ , with  $x_n \neq y_n$  and converging to  $p$ , such that for every CR function  $f \in \mathcal{O}(M)$  we have  $f(x_n) = f(y_n)$  for all  $n$ . Let  $d$  be the distance function on  $M$  defined by some Riemannian metric  $g$  on  $M$ . The functionals  $\xi_n$ , defined on a smooth function  $f$  on  $M$  by

$$(1.2) \quad \xi_n(f) = (f(x_n) - f(y_n)) / d(x_n, y_n),$$

converge, up to the choice of a subsequence, to a unit real tangent vector  $X \in T_pM$ . Clearly  $X(f) = 0$  for every  $f \in \mathcal{O}(M)$ , thus  $M$  is not strictly locally CR separable.  $\square$

Local strict CR separability is an open condition, because  $\dim S_x$  is upper semicontinuous with respect to  $x$ , and is actually equivalent to the existence, for each point  $x \in M$ , of a global CR map of  $M$  into a complex Euclidean space  $\mathbb{C}^n$  that is a CR embedding in a neighborhood of  $x$ .

We turn now to the case of parabolic CR manifolds. We briefly recall the construction of a parabolic CR manifold. For a thorough study of these CR manifolds we refer to [2].

Let  $G$  be a connected complex semisimple Lie group, and  $Q$  a parabolic subgroup of  $G$ . The homogeneous space  $Z = G/Q$  is a compact simply connected smooth complex projective variety, that is called a *flag manifold*. Fix a connected real form  $G_0$  of  $G$ . It acts on  $Z$  by left multiplication, partitioning  $Z$  into finitely many  $G_0$ -orbits. A *parabolic CR manifold*  $M$  is one of these  $G_0$  orbits.

If  $\mathfrak{g}$ ,  $\mathfrak{g}_0$  and  $\mathfrak{q}$  are the Lie algebras of  $G$ ,  $G_0$  and  $Q$  respectively, then the  $G_0$ -orbit  $G_0 \cdot \mathbf{o}$  through the point  $\mathbf{o} = eQ$  of  $Z$  is denoted by  $M(\mathfrak{g}_0, \mathfrak{q})$  and the pair  $(\mathfrak{g}_0, \mathfrak{q})$  is the *effective parabolic CR algebra* associated to  $M$  (see [6] for general definitions and properties of CR algebras). Since  $G$  acts transitively on  $Z$  we can always, up to conjugation of  $Q$  and  $G_0$  by an element of  $G$ , assume that  $\mathbf{o} \in M$ . Hence every  $G_0$ -orbit in  $Z$  is isomorphic to one of the form  $M = M(\mathfrak{g}_0, \mathfrak{q})$  for some choice of  $\mathfrak{g}_0$  and  $\mathfrak{q}$ .

Since  $G_0$  acts on  $Z$  by biholomorphisms,  $M$  inherits from  $Z$  a CR structure, and  $G_0$  acts on  $M$  by real analytic CR diffeomorphisms.

Let  $(\mathfrak{g}_0, \mathfrak{q})$  be an effective parabolic CR algebra, and  $M = M(\mathfrak{g}_0, \mathfrak{q})$  the associated parabolic CR manifold. Since  $\mathcal{S}$  is invariant under CR automorphisms of  $M$ , we have that  $\mathcal{S}$  is a  $G_0$ -homogeneous complex vector bundle on  $M$ . Furthermore we have (see [1, Lemma 14.5]):

**Lemma 1.2.** *The vector subspace of  $\mathfrak{g}$ :*

$$(1.3) \quad \mathfrak{s} = (\mathrm{d}\pi_e^{\mathbb{C}})^{-1}(S_{\mathbf{o}})$$

*is a parabolic complex Lie subalgebra of  $\mathfrak{g}$ , containing  $\mathfrak{q}$ .*  $\square$

Since  $\mathfrak{q} \subset \mathfrak{s}$ , we may consider the  $G_0$ -equivariant fibration:

$$(1.4) \quad \rho: M = M(\mathfrak{g}_0, \mathfrak{q}) \rightarrow M' = M(\mathfrak{g}_0, \mathfrak{s}).$$

Every CR function on  $M'$  defines, via the pullback by  $\rho$ , a CR function on  $M$ . Indeed more is true:

**Theorem 1.3.** *Let  $(\mathfrak{g}_0, \mathfrak{q})$  be an effective parabolic CR algebra and  $M = M(\mathfrak{g}_0, \mathfrak{q})$  the associated parabolic CR manifold. Then there exists a unique  $G_0$ -equivariant fibration  $\rho: M \rightarrow M'$  onto a strictly locally CR separable parabolic CR manifold  $M'$ , such that  $\rho$  induces an isomorphism on the space of CR functions, that is:*

$$(1.5) \quad \mathcal{O}(M) = \rho^* \mathcal{O}(M').$$

*The parabolic CR manifold  $M'$  is  $G_0$ -equivariantly CR isomorphic to  $M(\mathfrak{g}_0, \mathfrak{s})$ , where  $\mathfrak{s} = (\mathrm{d}\pi_e^{\mathbb{C}})^{-1}(S_{\mathbf{o}})$  is the Lie subalgebra of  $\mathfrak{g}$  defined in Lemma 1.2.*

*Proof.* From the definition of  $\mathfrak{s}$  it follows that  $M' = M(\mathfrak{g}_0, \mathfrak{s})$  has all the required properties, so we only need to prove uniqueness.

Let  $\rho'': M \rightarrow M'' = M(\mathfrak{g}_0, \mathfrak{q}'')$  be a  $G_0$ -equivariant fibration onto a strictly locally CR separable parabolic CR manifold  $M''$  such that  $\mathcal{O}(M) = \rho''^* \mathcal{O}(M'')$ . Then  $\mathfrak{q} \subset \mathfrak{q}''$  because  $\rho''$  is a  $G_0$ -equivariant fibration. On the other hand every CR function on  $M$  is the pullback of a CR function on  $M''$ , which is annihilated by  $T^{0,1}M''$ , hence  $\mathfrak{q}'' \subset \mathfrak{s}$ .

Finally  $S(M) \subset \rho^*(S(M''))$  because  $\rho$  is a CR map and  $S(M'') = T^{0,1}M''$  because  $M''$  is strictly CR separable. This yields  $\mathfrak{q}'' = \mathfrak{s}$ , as stated.  $\square$

We refer to the  $G_0$ -equivariant fibration (1.4), or to the corresponding  $\mathfrak{g}_0$ -equivariant fibration of CR algebras, as the *strictly CR separable reduction* of  $M(\mathfrak{g}_0, \mathfrak{q})$ , or of  $(\mathfrak{g}_0, \mathfrak{q})$ .

We can consider only simple parabolic CR manifolds. Indeed we have:

**Theorem 1.4.** *Let  $\mathfrak{g}_0 = \bigoplus_i \mathfrak{g}_{0_i}$  be the decomposition of the real semisimple Lie algebra  $\mathfrak{g}_0$  into the direct product of its simple ideals, and let  $\mathfrak{g}_i = \mathfrak{g}_{0_i}^{\mathbb{C}}$ ,  $\mathfrak{q}_i = \mathfrak{g}_i \cap \mathfrak{q}$ . Then each  $\mathfrak{q}_i$  is parabolic in  $\mathfrak{g}_i$ ,  $\mathfrak{q} = \bigoplus_i \mathfrak{q}_i$  and  $M(\mathfrak{g}_0, \mathfrak{q})$  is weakly (weakly locally, strictly locally) CR separable if and only if all  $M(\mathfrak{g}_{0_i}, \mathfrak{q}_i)$ 's are weakly (weakly locally, strictly locally) CR separable.*

*Proof.* The parabolic CR manifold  $M(\mathfrak{g}_0, \mathfrak{q})$  is isomorphic (see [9, § 5.4]) to the Cartesian product  $\prod_i M(\mathfrak{g}_{0_i}, \mathfrak{q}_i)$ .  $\square$

## 2. RESTRICTION TO MANIFOLDS OF FINITE TYPE

The following Theorem shows that we can restrict our consideration to parabolic CR manifolds of finite type.

**Theorem 2.1.** *Let  $M = M(\mathfrak{g}_0, \mathfrak{q})$  be a parabolic CR manifold, denote by  $M'$  the fiber and by  $M'' = M(\mathfrak{g}_0, \mathfrak{q}'')$  the base of a  $\mathbf{G}_0$ -equivariant CR fibration  $\phi: M \rightarrow M''$  onto a totally real parabolic CR manifold  $M''$ . Then  $M$  is weakly (weakly locally, strictly locally) CR separable if and only if  $M'$  is weakly (weakly locally, strictly locally) CR separable.*

First we prove a lemma:

**Lemma 2.2.** *Let  $M = M(\mathfrak{g}_0, \mathfrak{q})$  be a parabolic CR manifold,  $M'$  the fiber and  $M'' = M(\mathfrak{g}_0, \mathfrak{q}'')$  the base of a  $\mathbf{G}_0$ -equivariant CR fibration  $\phi: M \rightarrow M''$  onto a totally real parabolic CR manifold  $M''$ . Then every  $x \in M''$  has an open neighborhood  $U \subset M''$  such that  $\phi^{-1}(U)$  is CR diffeomorphic by a real analytic map to  $U \times M'$ .*

In particular Theorem 2.1 and Lemma 2.2 apply when  $\phi$  is the fundamental reduction [6, § 5] of  $M$ .

*Proof of Lemma 2.2.* Let  $\mathfrak{l}_0 = \mathfrak{g}_0 \cap \mathfrak{q}$  be the isotropy subalgebra of  $M$  and  $\mathfrak{l}_0'' = \mathfrak{g}_0 \cap \mathfrak{q}''$  that of  $M''$ . Since  $M''$  is totally real,  $\mathfrak{l}_0''$  is a parabolic subalgebra of  $\mathfrak{g}_0$ , hence there exists a nilpotent subalgebra  $\mathfrak{n}_0$  complementary to  $\mathfrak{l}_0''$ . Let  $\mathbf{L}_0, \mathbf{L}_0'', \mathbf{N}_0$  be the analytic subgroups of  $\mathbf{G}_0$  with Lie algebras  $\mathfrak{l}_0, \mathfrak{l}_0'', \mathfrak{n}_0$ , respectively and  $\pi: \mathbf{G}_0 \rightarrow M'' = \mathbf{G}_0/\mathbf{L}_0''$  the projection onto the quotient.

The restriction of  $\pi$  to  $\mathbf{N}_0$  is a real analytic local diffeomorphism. Choose an open neighborhood  $W$  of the identity in  $\mathbf{N}_0$  such that  $\pi|_W$  is a diffeomorphism onto an open subset  $\pi(W) = U$  of  $M''$ . Then the map:

$$(2.1) \quad \psi: U \times M' \ni (z, l\mathbf{L}_0) \mapsto ((\pi|_W)^{-1}(z)l)\mathbf{L}_0 \in M$$

is a real analytic CR trivialization in a neighborhood of  $e\mathbf{L}''$ . The result follows because of the homogeneity of  $M''$ .  $\square$

*Proof of Theorem 2.1.* Let  $x \neq y$  be two distinct points of  $M$ . If  $\phi(x) \neq \phi(y)$  then we can choose any function  $f$  on  $M''$  such that  $f \circ \phi(x) \neq f \circ \phi(y)$ , and  $f \circ \phi$  is CR, and separates  $x$  and  $y$ . If  $\phi(x) = \phi(y)$  then by Lemma 2.2 we can find a CR function  $f$  on  $M$  that separates  $x$  and  $y$  if and only if we can find such an  $f$  on  $\phi^{-1}(\phi(x))$ . Thus  $M$  is weakly (weakly locally) CR separable if and only if  $M'$  is weakly (weakly locally) CR separable.

Fix a point  $x \in M$ , let  $M' = \phi^{-1}(\phi(x))$  and denote by  $\iota: M' \rightarrow M$  the inclusion map. Let  $X \in T_x^{\mathbb{C}}M$  be a complex tangent vector at  $x$  with  $d\phi^{\mathbb{C}}(X) \neq 0$  and  $f$  a real analytic function on  $M''$  such that  $d\phi^{\mathbb{C}}(X)(f) \neq 0$ .

Then  $f \circ \phi$  is a CR function on  $M$  and  $X(f \circ \phi) \neq 0$ . This shows that  $\mathcal{S}(M) = \iota_*(\mathcal{S}(M'))$ , hence  $M$  is strictly locally CR separable if and only if  $M'$  is strictly locally CR separable.  $\square$

### 3. EXTENSION TO LEVI-FLAT ORBITS

The case of totally complex parabolic CR manifolds, was discussed by Wolf in [9]. There he proved (see [4, Thm. 4.4.3]) the following:

**Proposition 3.1.** *Let  $(\mathfrak{g}_0, \mathfrak{q})$  be a simple totally complex parabolic effective CR algebra and  $M = M(\mathfrak{g}_0, \mathfrak{q})$  the corresponding totally complex parabolic CR manifold. Then  $M$  is weakly locally CR separable if and only if  $M$  is a bounded symmetric domain. In this case  $M$  is also weakly CR separable and strictly locally CR separable.*  $\square$

We recall that a CR manifold  $M$  is *Levi-flat* if the analytic tangent distribution  $HM$  is integrable. For parabolic CR manifolds this is equivalent to the condition that the fibers of their fundamental reduction are totally complex. We have:

**Theorem 3.2.** *Let  $M = M(\mathfrak{g}_0, \mathfrak{q})$  be a  $G_0$ -orbit in the complex flag manifold  $Z = G/Q$ . Then there exists a Levi-flat  $G_0$ -orbit  $N$  in  $Z$ , with  $M \subset \overline{N}$ , such that every CR function  $f$  on  $M$  continuously extends to a function  $\tilde{f}$ , continuous on  $M \cup N$  and CR on  $N$ .*

*If  $M$  is of finite type, then  $N$  is totally complex, hence open in  $Z$ .*

*Proof.* If  $M$  is Levi-flat we take  $M' = M$ . Otherwise, a theorem of Tumanov [8] asserts that there exists a complex wedge  $W$ , with edge contained in  $M$ , such that every CR function  $f$  on  $M$  extends, continuously and uniquely, to a continuous function  $\tilde{f}$  on  $M \cup W$  that is holomorphic on  $W$ . Here by a complex wedge  $W$  with edge in  $M$  we mean a connected open subset  $W$  of a complex submanifold  $V$  of positive dimension of  $Z$  such that  $M \cap V$  is CR generic in  $V$  and contained in the closure  $\overline{W}$ .

Let  $x \in W$  and define, for all  $g \in G_0$ , a new  $\tilde{f}$  by setting:

$$(3.1) \quad \tilde{f}(g \cdot x) = (\widehat{f \circ m_g})(x),$$

where  $m_g: M \rightarrow M$  denotes the action of  $g$  on  $M$ . The function  $\tilde{f}$  is well defined and CR on the whole  $G_0$ -orbit  $M' = G_0 \cdot x$  through  $x$ . By choosing  $x$  close enough to  $M$ , we may arrange that  $M \subset \overline{M'}$  and  $\tilde{f}$  is continuous on  $M \cup M'$ .

By iterating this construction, we obtain a sequence of  $G_0$ -orbits  $M^{(i)}$  of nondecreasing dimension, each contained in the closure of the next. This sequence must necessary stabilize to a term  $M^{(k)} = N$ , that satisfies the first assertion of the theorem.

If  $M$  is of finite type, then also  $N$  is of finite type and, being Levi flat and CR generic, is open in  $Z$ .  $\square$

As a corollary, we obtain:

**Corollary 3.3.** *If  $M = M(\mathfrak{g}_0, \mathfrak{q})$  is a strictly locally CR separable parabolic CR manifold, embedded in the complex flag manifold  $Z = G/Q$ , then:*

- (1) *there exists a strictly locally CR separable Levi-flat  $G_0$ -orbit  $N \subset Z$  with  $M \subset \overline{N}$ ;*
- (2)  *$M$  is (globally) weakly CR separable.*

*If  $M$  is of finite type then  $N$  is a bounded symmetric domain.*

*Proof.* We may assume that  $M$  is the  $G_0$ -orbit in  $Z$  through the point  $\mathfrak{o} = eQ$ .

Let  $N$  be the Levi-flat  $G_0$ -orbit defined in Theorem 3.2. Let  $\phi: Z \rightarrow Z' = G/Q'$  be the  $G$ -equivariant fibration of complex flag manifolds that induces, by restriction to  $N$ , the strictly separable CR reduction  $\phi|_N: N \rightarrow N' \subset Z'$ . Then every CR function  $f$  on  $M$  extends continuously to a function  $\tilde{f}$ , continuous on  $M \cup N$  and CR on  $N$ , constant along the fibers of  $\phi$ . By continuity also  $f$  is constant along the fibers of  $\phi$  and furthermore  $f = \phi \circ f'$  for some CR function  $f'$  on  $M' = \phi(M)$ . This shows that:

$$(3.2) \quad S_{\mathfrak{o}}(M) \supset (d\phi^{\mathbb{C}})^{-1} T_{\mathfrak{o}}^{(0,1)} M'.$$

Since  $M$  is strictly CR separable, we obtain that:

$$(3.3) \quad T_{\mathfrak{o}}^{(0,1)} M = (d\phi^{\mathbb{C}})^{-1} T_{\mathfrak{o}}^{(0,1)} M',$$

which in turn implies that  $\mathfrak{q} = \mathfrak{q}'$ . Thus  $N = N'$ , that is  $N$  is strictly locally CR separable.

If  $M$  is of finite type, by Proposition 3.1,  $N$  is a bounded symmetric domain and  $M \subset \overline{N}$ . This fact also implies that  $M$  is weakly CR separable, thus the Theorem is proved if  $M$  is of finite type.

If  $M$  is not of finite type, we apply two times Theorem 2.1 to the fiber  $M'$  of its fundamental reduction and obtain:

$$\begin{aligned} M \text{ is strictly locally CR separable} &\implies \\ &\implies M' \text{ is strictly locally CR separable} \implies \\ &\implies M' \text{ is weakly CR separable} \implies \\ &\implies M \text{ is weakly CR separable,} \end{aligned}$$

completing the proof.  $\square$

#### 4. EXAMPLES

In this section we discuss some examples. We will not need to utilize Tumanov's results, but more elementary extension theorems will suffice.

In particular we recall the following statement. Let  $S^3 = \{z \in \mathbb{C}^2 \mid |z| = 1\}$  be the three-dimensional sphere, endowed with the usual CR structure,  $B^2 = \{z \in \mathbb{C}^2 \mid |z| < 1\}$  the two-dimensional complex ball,  $\Sigma$  a real two dimensional linear subspace of  $\mathbb{C}^2$  (that may or may not be a complex line) and set:  $\check{S}^3 = S^3 \setminus \Sigma$ . Then every CR function on  $\check{S}^3$  extends continuously to a function continuous on  $\overline{B}^2 \setminus \Sigma$  and holomorphic on  $\check{B}^2 = B^2 \setminus \Sigma$ .

For totally complex parabolic CR manifolds and minimal parabolic CR manifolds the obstruction to CR separability is the existence of embedded compact complex submanifold. The general case is quite different. Indeed we exhibit two examples of parabolic CR manifold that are not weakly locally CR separable, but do not contain any compact complex submanifold.

**Example 1.** Let  $H(u, v) = u^*Av$  be the Hermitian form on  $\mathbb{C}^3$  associated to the matrix  $A = \text{diag}(-1, 1, 1)$ , and  $G \simeq \text{SL}(3, \mathbb{C})$  the group of unimodular complex matrices. The subgroup  $G_0$  of matrices in  $G$  that leave  $H$  invariant is a real form of  $G$ , isomorphic to  $\text{SU}(1, 2)$ .

Let  $Z$  be the complex flag manifold:

$$(4.1) \quad Z = \{\ell_1 \subset \ell_2 \subset \mathbb{C}^3 \mid \dim \ell_i = i\}$$

and  $M$  the parabolic CR manifold:

$$(4.2) \quad M = \{(\ell_1, \ell_2) \in Z \mid \ell_1 \text{ is } H\text{-isotropic}, \ell_2 \text{ is } H\text{-hyperbolic}\}.$$

Consider the  $G$ -equivariant fibration

$$(4.3) \quad \phi: Z \ni (\ell_1, \ell_2) \mapsto \ell_1 \in W$$

onto the complex flag manifold:

$$(4.4) \quad W = \{\ell_1 \subset \mathbb{C}^3 \mid \dim \ell_1 = 1\}.$$

Then  $\phi$  restricts to a  $G_0$ -equivariant fibration  $\phi: M \rightarrow N$  onto the parabolic CR manifold:

$$(4.5) \quad N = \{\ell_1 \in W \mid \ell_1 \text{ is } H\text{-isotropic}\}.$$

The fiber of  $\phi|_M$  over a point  $\ell_1 \in N$  is the set  $\{\ell_2 \subset \mathbb{C}^3 \mid \ell_1 \subset \ell_2 \not\subset (\ell_1)^\perp\}$ , which is biholomorphic to  $\mathbb{C}$ . The CR manifold  $N$  is the boundary of the open domain:

$$(4.6) \quad D = \{\ell_1 \in W \mid \ell_1 \text{ is } H\text{-negative}\}.$$

Fix an  $H$ -positive line  $\hat{\ell}_1^+ \subset \mathbb{C}^3$  and define:

$$(4.7) \quad M_{\hat{\ell}_1^+} = \{(\ell_1, \ell_2) \in M \mid \ell_2 = \ell_1 + \hat{\ell}_1^+\} \simeq \check{S}^3.$$

Any CR function  $f$  on  $M_{\hat{\ell}_1^+}$  extends continuously to a function  $\tilde{f}$ , continuous on  $M_{\hat{\ell}_1^+} \cup U_{\hat{\ell}_1^+}$  and holomorphic on  $U_{\hat{\ell}_1^+}$ , where:

$$(4.8) \quad U_{\hat{\ell}_1^+} = \{(\ell_1, \ell_2) \in Z \mid \ell_1 \text{ is } H\text{-negative}, \ell_2 = \ell_1 + \hat{\ell}_1^+\} \simeq \check{B}^2.$$

By letting  $\hat{\ell}_1^+$  vary among all  $H$ -positive lines, we obtain that every CR function  $f$  on  $M$  extends continuously to a function  $\tilde{f}$  continuous on  $M \cup U$  and holomorphic on  $U$ , where:

$$(4.9) \quad U = \{(\ell_1, \ell_2) \in Z \mid \ell_1 \text{ is } H\text{-negative and } \ell_2 \text{ is } H\text{-hyperbolic}\}$$

Let:

$$(4.10) \quad \begin{aligned} V &= \phi^{-1}(D) \setminus U \\ &= \{(\ell_1, \ell_2) \in Z \mid \ell_1 \text{ is } H\text{-negative}, H \text{ has signature } (0, -) \text{ on } \ell_2\}. \end{aligned}$$

Then  $V$  has real codimension two in  $\phi^{-1}(D)$  and is not complex analytic. Hence, by a theorem of Hartogs ([5], see [7, § 4, Thm. 3]), there is a point  $x \in V$  with the property that  $\tilde{f}$  holomorphically extends to a full neighborhood  $U_x$  of  $x$  in  $Z$ . It follows that  $\tilde{f}$  is constant on  $\phi^{-1} \circ \phi(y)$  for all  $y \in U_x$ , and by unique continuation  $\tilde{f}$  is constant along the fibers of  $\phi$ , hence  $M$  is not weakly locally CR separable.

**Example 2.** Let  $B(u, v) = {}^t u A v$  and  $H(u, v) = u^* A v$  be the bilinear and the Hermitian form on  $\mathbb{C}^5$ , associated to the matrix  $A = \text{diag}(-1, -1, 1, 1, 1)$ , and  $G \simeq \text{SO}(5, \mathbb{C})$  the group of unimodular complex matrices that preserve  $B$ . The connected component  $G_0$  of the identity in the subgroup of the real matrices of  $G$  is isomorphic to  $\text{SO}^0(2, 3)$  and the elements of  $G_0$  also preserve the Hermitian form  $H$ .

Let  $Z$  be the complex flag manifold:

$$(4.11) \quad Z = \{\ell_1 \subset \ell_2 \subset \mathbb{C}^5 \mid \dim \ell_i = i, \ell_i \text{ is } B\text{-isotropic}\}$$

and  $M$  the parabolic CR manifold:

$$(4.12) \quad M = \{(\ell_1, \ell_2) \in Z \mid \ell_1 \text{ is } H\text{-isotropic}, \ell_1 \neq \bar{\ell}_1, \ell_2 \text{ is } H\text{-hyperbolic}\}.$$

Consider the  $G$ -equivariant fibration

$$(4.13) \quad \phi: Z \ni (\ell_1, \ell_2) \mapsto \ell_1 \in W$$

onto the complex flag manifold:

$$(4.14) \quad W = \{\ell_1 \subset \mathbb{C}^5 \mid \dim \ell_1 = 1, \ell_1 \text{ is } B\text{-isotropic}\}.$$

Then  $\phi$  restricts to a  $G_0$ -equivariant fibration  $\phi: M \rightarrow N$  onto the parabolic CR manifold:

$$(4.15) \quad N = \{\ell_1 \in W \mid \ell_1 \text{ is } H\text{-isotropic}, \ell_1 \neq \bar{\ell}_1\}.$$

The CR manifold  $N$  is an open stratum in the boundary of the open domain:

$$(4.16) \quad D = \{\ell_1 \in W \mid \ell_1 \text{ is } H\text{-negative}\}.$$

Fix a  $B$ -isotropic and  $H$ -positive line  $\hat{\ell}_1^+ \subset \mathbb{C}^5$  and define:

$$(4.17) \quad M_{\hat{\ell}_1^+} = \{(\ell_1, \ell_2) \in M \mid \ell_2 = \ell_1 + \hat{\ell}_1^+\} \simeq \check{S}^3.$$

Any CR function on  $M_{\hat{\ell}_1^+}$  continuously extends to a function  $\tilde{f}$ , continuous on  $M_{\hat{\ell}_1^+} \cup U_{\hat{\ell}_1^+}$  and holomorphic on  $U_{\hat{\ell}_1^+}$ , where:

$$(4.18) \quad U_{\hat{\ell}_1^+} = \{(\ell_1, \ell_2) \in Z \mid \ell_1 \text{ is } H\text{-negative}, \ell_2 = \ell_1 + \hat{\ell}_1^+\} \simeq \check{B}^2.$$

By letting  $\hat{\ell}_1^+$  vary among all  $B$ -isotropic and  $H$ -positive lines, we obtain that every CR function  $f$  on  $M$  continuously extends to a function  $\tilde{f}$ , continuous on  $M \cup U$  and holomorphic on  $U$ , where:

$$(4.19) \quad U = \{(\ell_1, \ell_2) \in Z \mid \ell_1 \text{ is } H\text{-negative and } \ell_2 \text{ is } H\text{-hyperbolic}\}.$$

Let:

$$(4.20) \quad \begin{aligned} V &= \phi^{-1}(D) \setminus U \\ &= \{(\ell_1, \ell_2) \in Z \mid \ell_1 \text{ is } H\text{-negative and } \ell_2 \text{ is } H\text{-degenerate}\}. \end{aligned}$$

Then  $V$  has real codimension two in  $\phi^{-1}(D)$  and is not complex analytic, hence by using the same argument of Example 1, it follows that any CR function on  $M$  is constant along the fibers of  $\phi$ . Thus  $M$  is not weakly locally CR separable.

The next example consists of a parabolic CR manifold that is weakly, but not strictly, locally CR separable.



**Example 3.** Let  $G \simeq \mathrm{SL}(4, \mathbb{C})$  be the group of unimodular  $4 \times 4$  complex matrices and  $G_0 \simeq \mathrm{SU}(2, 2)$  the subgroup of matrices preserving the Hermitian form  $H$  associated to the matrix  $\mathrm{diag}(-1, -1, 1, 1)$ .

Let  $Z$  be the complex flag manifold:

$$(4.21) \quad Z = \{\ell_1 \subset \ell_2 \subset \mathbb{C}^5 \mid \dim \ell_i = i\},$$

and  $M$  the parabolic CR manifold:

$$(4.22) \quad M = \{(\ell_1, \ell_2) \in Z \mid \ell_1 \text{ is } H\text{-isotropic, } H|_{\ell_2} \text{ has signature } (0, -)\}.$$

Consider the  $G$ -equivariant fibration

$$(4.23) \quad \phi: Z \ni (\ell_1, \ell_2) \mapsto \ell_2 \in W$$

onto the complex flag manifold:

$$(4.24) \quad W = \{\ell_2 \subset \mathbb{C}^5 \mid \dim \ell_2 = 2\}.$$

Then  $\phi$  restricts to a  $G_0$ -equivariant fibration  $\phi: M \rightarrow N$ , which is a CR map and a smooth diffeomorphism, but not a CR fibration, onto the parabolic CR manifold:

$$(4.25) \quad N = \{\ell_2 \in W \mid H|_{\ell_2} \text{ has signature } (0, -)\}.$$

The CR manifold  $N$  is an open stratum in the boundary of the open domain:

$$(4.26) \quad D = \{\ell_2 \in W \mid \ell_2 \text{ is } H\text{-negative}\}.$$

Fix an  $H$ -negative line  $\hat{\ell}_1^- \subset \mathbb{C}^5$  and define:

$$(4.27) \quad M_{\hat{\ell}_1^-} = \{(\ell_1, \ell_2) \in M \mid \ell_1 \text{ is } H\text{-isotropic, } \ell_1 \perp_H \hat{\ell}_1^-, \ell_2 = \ell_1 + \hat{\ell}_1^-\} \simeq S^3.$$

Any CR function on  $M_{\hat{\ell}_1^-}$  continuously extends to a function continuous on  $M_{\hat{\ell}_1^-} \cup U_{\hat{\ell}_1^-}$ , and holomorphic on  $U_{\hat{\ell}_1^-}$ , where:

$$(4.28) \quad U_{\hat{\ell}_1^-} = \{(\ell_1, \ell_2) \in Z \mid \ell_1 \text{ is } H\text{-negative, } \ell_1 \perp_H \hat{\ell}_1^-, \ell_2 = \ell_1 + \hat{\ell}_1^-\} \simeq B^2.$$

By letting  $\hat{\ell}_1^-$  vary among all  $B$ -isotropic and  $H$ -positive lines, we obtain that every CR function on  $M$  continuously extends to a function continuous on  $M \cup U$  and holomorphic on  $U$ , where:

$$(4.29) \quad U = \{(\ell_1, \ell_2) \in Z \mid \ell_2 \text{ is } H\text{-negative}\}.$$

Since each fiber of the restriction of  $\phi$  to  $U$  is biholomorphic to  $\mathbb{CP}^1$ , every CR function  $f$  on  $M$  can be extended to a CR function  $\tilde{f}$  on  $\phi^{-1}(N)$ , which is constant along the fibers of  $\phi$ . This shows that  $f$  is also CR on  $N$ , hence  $M$  is not strictly locally CR separable.

On the other hand  $N$  is strictly locally CR separable, hence by Lemma 1.1  $M$  is weakly locally CR separable.

## REFERENCES

1. Andrea Altomani, Costantino Medori, and Mauro Nacinovich, *The CR structure of minimal orbits in complex flag manifolds*, J. Lie Theory **16** (2006), no. 3, 483–530. MR 2248142
2. Andrea Altomani, Costantino Medori, and Mauro Nacinovich, *Orbits of real forms in complex flag manifolds*, preprint, Nov 2006, arXiv:math.CV/0611755.

3. M. Salah Baouendi, Peter Ebenfelt, and Linda Preiss Rothschild, *Real submanifolds in complex space and their mappings*, Princeton Mathematical Series, vol. 47, Princeton University Press, Princeton, NJ, 1999. MR 1668103
4. Gregor Fels, Alan Huckleberry, and Joseph A. Wolf, *Cycle spaces of flag domains*, Progress in Mathematics, vol. 245, Birkhäuser Boston Inc., Boston, MA, 2006, A complex geometric viewpoint. MR 2188135
5. Friedrich Hartogs, *Über die aus den singulären Stellen einer analytischen Funktion mehrerer Veränderlichen bestehenden Gebilde.*, Acta Math. **32** (1909), 57–79.
6. Costantino Medori and Mauro Nacinovich, *Algebras of infinitesimal CR automorphisms*, J. Algebra **287** (2005), no. 1, 234–274. MR 2134266
7. Raghavan Narasimhan, *Several complex variables*, The University of Chicago Press, Chicago-London, 1971, Chicago Lectures in Mathematics. MR 0342725
8. Alexander E. Tumanov, *Extension of CR-functions into a wedge*, Mat. Sb. **181** (1990), no. 7, 951–964. MR 1070489
9. Joseph A. Wolf, *The action of a real semisimple group on a complex flag manifold. I. Orbit structure and holomorphic arc components*, Bull. Amer. Math. Soc. **75** (1969), 1121–1237. MR 0251246

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI ROMA “TOR VERGATA”, VIA DELLA RICERCA SCIENTIFICA, 00133 ROMA (ITALY)

*E-mail address:* altomani@mat.uniroma2.it